


# Flaring Activity in a Black Hole Magnetosphere

Enzo Figueiredo


Benoît Cerutti, John Mehlhaff, Adrien Soudais




European  
Research  
Council



# Black holes and jet emission




Evidences for a connection between SMBH and galactic jets  
Non thermal emission → Particle acceleration




No jet? But flares!

# Sagittarius A\*: a not-so-quiet black hole

## GRAVITY observations:




- Time-duration: 30–90 min
- Near-infrared + X-rays
- Close to BH ( $6 r_g$ )




- Large-scale magnetic field (10–15 G)
- Orientation orthogonal to rotation axis

# Sagittarius A\*: about its environment

Wind-fed accretion from surrounding stars in the Galactic Center




Ressler+2020



Large-scale simulations predict a low-angular momentum plasma that is highly magnetized! ( $\beta = \text{few}$ )

# GRZeltron: a GRPIC code (Parfrey+2019)

3+1 formalism (Komissarov, 2004)



Field Interpolation

Particle Push

$$\dot{x}^i = \frac{\alpha}{\Gamma} \gamma^{ij} u_j - \beta^i$$


$$\dot{u}^i = \text{Lorentz} + \text{Metric}$$

Fields Evolution

$$\frac{\partial B}{\partial t} = -c \nabla \times E$$

$$\frac{\partial D}{\partial t} = c \nabla \times H - 4\pi \mathbf{J}$$


Kerr metric, KS spherical coordinates



Charge/Current Deposition

- Full treatment of non ideal phenomena
- Can capture microphysics (e.g. QED processes)
- Acceleration of particles self consistently treated
- Straightforward radiative transfer

# Numerical setup



$30 r_g$

-Schwarzschild black hole

-Axisymmetric setup

-Plasma injected on the outer-edge of the box (free-fall)


-Pair plasma at “low” magnetization

$$\sigma_0 = \frac{B_0^2}{4\pi n_0 m_e c^2} = 0.1$$


 Injection area

 Magnetic field

# Flux eruptions in the black hole surroundings




# Strong particle acceleration at the eruptions





$\sim \sigma$  near the event horizon!

The eruptive phase should correspond to flare of synchrotron emission




Typical timescale:  $100 t_g$   
 $\rightarrow 30$  min for Sgr A\* mass

# What if we put ions?



Extrapolating to the real mass ratio:  $\langle \gamma_e \rangle \sim 10^4$

# Conclusions



- A simple setup can capture flaring activity of a black hole magnetosphere from first principles
- The universality of magnetic reconnection appears to have a central role to explain the eruptions timescales

Incoming 3D simulations!

