

DUST EVOLUTION MODELING IN THE DISK-BUILDING STAGE



Maxime Lombart (maxime.lombart@cea.fr)
postdoc in ERC project PEBBLES/CEA Paris-Saclay

Wada et al. 2018

With: U. Lebreuilly, A. Maury


Collaborators: G. Verrier, C.-E.
Bréhier, M. Hutchison, Y.-N.
Lee, G. Laibe



ASNUM, 2025/12/17, Grenoble

STAR FORMATION, DISK EVOLUTION AND PLANET FORMATION 2


The first seeds for planets may form in less than a few million years.
How early do pebbles form around YSOs ?




Dust growth must be an efficient process !

3D SIMULATIONS WITH DUST COAGULATION/FRAGMENTATION 3



discs




Vericel et al., 2021



protostellar
collapses



Tsukamoto et al., 2021



gas and dust dynamic in 1D

single dust size approximation

dust size distribution

gas and dust dynamic in 3D



3D SIMULATIONS WITH DUST COAGULATION

4


Marian Smoluchowski (1872-1917)



Model:

- mean-field theory (fluctuation of physical quantities are neglected)
- binary collision
- spherical particles
- collision with coalescence






m, m' : masses

n : number density of particles per mass unit

K : collision kernel



$$\frac{\partial n(m, t)}{\partial t} = \underbrace{\frac{1}{2} \int_0^m K(m', m - m') n(m', t) n(m - m', t) dm'}_{\text{gain}} - \underbrace{n(m, t) \int_0^\infty K(m, m') n(m', t) dm'}_{\text{loss}}$$


gain

loss

$$K(m_1, m_2) = \sigma(m_1, m_2) \Delta v(m_1, m_2) \rightarrow \text{need numerical solutions}$$

Requirements

- ◇ 3 orders of magnitude in size ($1\mu\text{m} - 1\text{mm}$) $\Leftrightarrow$ 9 orders in mass
- ◇ mass range discretization $\rightarrow$ 20 mass bins (low numerical cost)



Discontinuous Galerkin scheme


conservative form (Tanaka et al., 1996)

Liu et al., 2019
Lombart &
Laibe, 2021

$$\begin{cases} \frac{\partial g(m, t)}{\partial t} + \frac{\partial F_{\text{coag}}[g](m, t)}{\partial m} = 0 \\ F_{\text{coag}}[g](m, t) = \int_0^m \int_{m-m_1}^{\infty} \frac{K(m_1, m_2)}{m_2} g(m_1, t) g(m_2, t) dm_2 dm_1 \end{cases}$$



COAgulation with
Lightning fast
Algorithm




→ high accuracy
→ few number of mass bins

BENCHMARK COALA

7

Lombart & Laibe (2021), Laibe & Lombart (2022)



Requirements

- ✓ 9 orders of magnitude in mass
- ✓ 10-20 dust size bins

3D simulations with
polydisperse coagulation



DUST GROWTH IN PROTOSTELLAR COLLAPSE

8

Lombart et al., 2025 (minor revisions)

RAMSES MHD model of B335-like
protostar
gas & dust (40 bins)

Implementation:

- dust growth as source term of continuity equation


$$\frac{\partial \rho_k}{\partial t} + \nabla \cdot [\rho_k \mathbf{v}_k] = \left. \frac{\partial \rho_k}{\partial t} \right|_{\text{growth}}$$

$$\rho_k = \int_{I_k} g(m, t) dm$$


- ordinary operator splitting

$$\rho_k^* = \rho_k(t_n) - \Delta t \left(\nabla \cdot [\rho_k \mathbf{v}_k] \right) \rightarrow \text{RAMSES}$$

$$\rho_k(t_{n+1}) = \rho_k^* + \Delta t \left. \frac{\partial \rho_k}{\partial t} \right|_{\text{growth}} \rightarrow \text{COALA}$$






See Gabriel Verrier's talk
for multi-fluid RAMSES



DUST GROWTH IN PROTOSTELLAR COLLAPSE

Lombart et al., 2025 (minor revisions)



Mass of grains in the envelope
(in % of total dust mass)

Initially (MRN distribution):

- $\sim 0,25 \mu\text{m} \rightarrow 20\%$
- $\sim 5 \mu\text{m} \rightarrow 0\%$

After 100 kyrs:

- $\sim 0,25 \mu\text{m} \rightarrow 13\%$
- $\sim 5 \mu\text{m} \rightarrow 1\%$

DUST GROWTH IN PROTOSTELLAR COLLAPSE

11

Lombart et al., 2025 (minor revisions)

Initial dust-to-gas ratio $\theta_0 = 0.01$

$$\epsilon = \frac{\rho_d}{\rho_g + \rho_d} \text{ and } \epsilon_0 = \frac{\theta_0}{1 + \theta_0}$$


Main mechanism for differential grain-grain velocity is from turbulence
Ormel's model:

$$\Delta v_{i,j} \propto \sqrt{\alpha} c_s \sqrt{S_t}$$


$\alpha \rightarrow$ level of turbulence

In envelope: $\alpha = 1.5$


In Disk ($n_g > 10^9 \text{cm}^{-3}$): $\alpha = 10^{-4}$



Lombart et al., 2025 (minor revisions)



DUST FRAGMENTATION MODEL




Safronov, 1972; Jones et al., 1996; Suttner et al., 2001; Blum, 2006; Hirashita et al., 2009; Rafikov et al., 2020

Conservative form (Lombart et al. 2024):

$$F_{\text{frag}}[g](m, t) = \frac{1}{2} \int_0^m \int_0^\infty \int_0^\infty \mathbf{1}_{m_1+m_2 \geq m} \frac{m_f}{m_1 m_2} b(m_f, m_1, m_2) K(m_1, m_2) g(m_1, t) g(m_2, t) dm_2 dm_1 dm_f$$
$$- \int_0^m \int_{m-m_1}^\infty \frac{K(m_1, m_2)}{m_2} g(m_1, t) g(m_2, t) dm_2 dm_1$$

Bukhari Syed et al. 2017



DUST FRAGMENTATION MODEL



Benchmark for fragmentation

Feingold et al., 1988


$$K(u, v) = 1$$

$$b(w, u, v) = \gamma^2(u + v)e^{-\gamma w}$$


3D simulations
with polydisperse
general non-linear
fragmentation




Lombart et al. 2024



Growth and fragmentation of multi-component dust aggregates





Need 2D Smoluchowski coagulation equation

$$\frac{\partial n(m_s, m_i, t)}{\partial t} = \frac{1}{2} \int_0^{m_s} \int_0^{m_i} K(m_s - m'_s, m'_s, m_i - m'_i, m'_i) n(m'_s, m'_i, t) n(m_s - m'_s, m_i - m'_i, t) dm'_s dm'_i - \int_0^\infty \int_0^\infty K(m_s, m'_s, m_i, m'_i) n(m_s, m_i, t) n(m'_s, m'_i, t) dm'_s dm'_i$$

$$K(m_s - m'_s, m'_s, m_i - m'_i, m'_i) = P_{\text{stick}}(m_s, m'_s, m_i, m'_i, \Delta v) \sigma \Delta v$$

Sticking probability

$$P_{\text{stick}}(m_s, m'_s, m_i, m'_i) = \epsilon_{s-s} \frac{m_{s,A}}{m_A} \frac{m_{s,B}}{m_B} + \epsilon_{s-i} \left(\frac{m_{s,A}}{m_A} \frac{m_{i,A}}{m_A} + \frac{m_{s,B}}{m_B} \frac{m_{i,B}}{m_B} \right) + \epsilon_{i-i} \frac{m_{i,A}}{m_A} \frac{m_{i,B}}{m_B}$$


Sticking efficiency of s-s collision

Sticking efficiency of s-i collision

Sticking efficiency of i-i collision

Need data from lab experiments

Dust aggregate is defined by its mass m and porosity $\psi = \frac{V}{V^*}$



Need 2D Smoluchowski coagulation equation

$$\frac{\partial n(m, \psi_m, \tau)}{\partial t} = \frac{1}{2} \int_0^m \int_{\psi_{\min}}^{\psi_{\max}} \int_{\psi_{\min}}^{\psi_{\max}} K(m - m_1, \psi_{m-m_1}; m_1, \psi_{m_1}) n(m_1, \psi_{m_1}, t) n(m - m_1, \psi_{m-m_1}, t) \delta(\psi_m - \Gamma(m - m_1, \psi_{m-m_1}; m_1, \psi_{m_1})) d\psi_{m-m_1} d\psi_{m_1} dm_1$$

$$- n(m, \psi_m, t) \int_0^\infty \int_{\psi_{\min}}^{\psi_{\max}} K(m, \psi_m; m_1, \psi_{m_1}) n(m_1, \psi_{m_1}, t) d\psi_{m_1} dm_1$$

$\Gamma(m - m_1, \psi_{m-m_1}; m_1, \psi_{m_1}) \rightarrow$ determine the porosity of the resulting aggregate




Multi-component dust:

→ how to model the composition of fragments ?

Dust aggregates:

→ how to model the mass and porosity of the fragments ?



Need data from lab experiments



Blum, 2006; Ormel et al., 2007; Okuzumi et al., 2009; Hirashita et al., 2021; Hasegawa et al., 2022




CHARACTERIZING DUST EVOLUTION DURING THE DISK-BUILDING STAGE 19



+



$\log_{10}(\alpha_{\text{peak}}) (\mu\text{m})$
-0.5 0.1 0.7 1.4 2.0



Carpine et al., 2025

Realistic dust
optical properties




THEMIS

The Heterogeneous dust Evolution Model for
Interstellar Solids



amorphous silicate core
+ carbon mantle

amorphous carbon
hydrogenated

Monomers
 $\sim 0.1 \mu\text{m}$



$\sim 1 \mu\text{m}$



aggregates

compact sphere

Radiative transfer
codes

Realistic synthetic
observations

- ◉ The size evolution of dust particles are modeled by Smoluchowski-like equations.
- ◉ **3D simulations with polydisperse dust coagulation/fragmentation**
 - ◉ 9 orders of magnitude in mass ($1\mu\text{m}$ - 1mm)
 - ◉ Few number of bins ~ 20
 - ◉ Good accuracy
- ◉ **Dust grains are predicted to grow by 2 orders of magnitude in collapsing envelopes (just gravity/turbulence, without magnetic effects yet)**
- ◉ Coagulation/fragmentation of dust aggregates and multicomponent dust can be modeled by **2D extension**.
- ◉ **PEBBLES**: realistic synthetic observations by combining simulations, radiative transfer with realistic dust optical properties

